

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
26 June 2003 (26.06.2003)

PCT

(10) International Publication Number
WO 03/051851 A1(51) International Patent Classification: C07D 241/24,
A61K 31/495, A61P 25/18S-431 83 (SE). GREASLEY, Peter [GB/SE]; Molndal,
S-431 83 (SE). TERRICABRA, Emma [ES/SE]; Molndal,
S-431 83 (SE). WILSTERMANN, Johan, Michael
[SE/SE]; Molndal, S-431 83 (SE).

(21) International Application Number: PCT/GB02/05742

(22) International Filing Date:
18 December 2002 (18.12.2002)(74) Agent: ASTRAZENECA; Global Intellectual Property,
Mereside, Alderley Park, Macclesfield, Cheshire SK10
4TG (GB).

(25) Filing Language: English

(26) Publication Language: English


(30) Priority Data:
0104330-6 19 December 2001 (19.12.2001) SE(71) Applicant (for all designated States except MG, US): AS-
TRAZENECA AB [SE/SE]; Sodertalje, S-151 85 (SE).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.(71) Applicant (for MG only): ASTRAZENECA UK LIM-
ITED [GB/GB]; 15 Stanhope Gate, London, Greater Lon-
don W1K 1LN (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BERGGREN,
Anna, Ingrid, Kristina [SE/SE]; Molndal, S-431 83
(SE). BOSTROM, Stig, Jonas [SE/SE]; Molndal, S-431
83 (SE). ELEBRING, Stig, Thomas [SE/SE]; Molndal,(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).Published:
— with international search report

[Continued on next page]

(54) Title: 5, 6-DIARYL-PYRAZINE-2-AMIDE DERIVATIVES AS CB1 ANTAGONISTS



(57) Abstract: The present invention relates to compounds of formula (I), and pharmaceutically acceptable salts, prodrugs, solvates and crystalline forms thereof, in which R¹ and R² independently represent: a C₁₋₆alkyl group; an optionally substituted (amino)C₁₋₄alkyl group; an optionally substituted non-aromatic C₃₋₁₅carbocyclic group; a (C₃₋₁₂cycloalkyl)C₁₋₃alkyl group; a group -(CH₂)_r(phenyl), in which r is 0, 1, 2, 3 or 4, s is 1 when r is 0 otherwise s is 1 or 2 and the phenyl groups are optionally independently substituted by Z; naphthyl; anthracenyl; an optionally substituted saturated 5 to 8 membered heterocyclic group containing one nitrogen and optionally one of the following: oxygen, sulphur or an additional nitrogen; 1-adamantylmethyl; a group -(CH₂)_tHet in which t is 0, 1, 2, 3 or 4, and the alkylene chain is optionally substituted and Het represents an optionally substituted aromatic heterocycle; or R¹ represents H and R² is as defined above; or R¹ and R² together with the nitrogen atom to which they are attached represent a saturated optionally substituted 5 to 8 membered heterocyclic group as defined above; X is CO or SO₂; Y is absent or represents NH optionally substituted by a C₁₋₃alkyl group; R³ and R⁴ independently represent phenyl, thienyl or pyridyl substituted by Z; Z represents a C₁₋₃alkyl group, a C₁₋₃alkoxy group, hydroxy, halo, trifluoromethyl, trifluoromethylthio, trifluoromethoxy, trifluoromethylsulphonyl, nitro, amino, mono or di C₁₋₃alkylamino, mono or di C₁₋₃alkylamido, C₁₋₃alkylsulphonyl, C₁₋₃alkoxycarbonyl, carboxy, cyano, carbamoyl, mono or di C₁₋₃alkyl carbamoyl, sulphonyl and acetyl; and R⁵ is H, a C₁₋₃alkyl group, a C₁₋₃alkoxymethyl group, trifluoromethyl, a hydroxyC₁₋₃alkyl group, C₁₋₃alkoxycarbonyl, carboxy, cyano, carbamoyl, mono or di C₁₋₃alkyl carbamoyl, acetyl, or hydrazinocarbonyl of formula -CONHNHNR⁶; with the provisos; and processes for preparing such compounds, their use in the treatment of obesity, psychiatric and neurological disorders, to methods for their therapeutic use and to pharmaceutical compositions containing them.



— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

5, 6-DIARYL-PYRAZINE-2-AMIDE DERIVATIVES AS CBI ANTAGONISTS

Field of invention

5 The present invention relates to certain pyrazine carboxamide compounds of formula I, to processes for preparing such compounds, to their use in the treatment of obesity, psychiatric and neurological disorders, to methods for their therapeutic use and to pharmaceutical compositions containing them.


10 Background of the invention

It is known that certain CB₁ modulators (known as antagonists or inverse agonists) are useful in the treatment of obesity, psychiatric and neurological disorders (WO01/70700 and EP 656354). However, there is a need for CB₁ modulators with improved
15 physicochemical properties and/or DMPK properties and/or pharmacodynamic properties.

Pyrazinecarboxamides are reported to possess antithrombotic properties (WO 92/ 02513). The compounds disclosed in this document are disclaimed from the compound claims of the present invention. 5,6-Diphenyl-2-pyrazinecarboxylic acid is disclosed in CH 458 361.
20

Description of the invention

The invention relates to compounds of the general formula (I)
25



I

and pharmaceutically acceptable salts, prodrugs, solvates and crystalline forms thereof, in which

R^1 and R^2 independently represent :

a C_{1-6} alkyl group;

5 an (amino) C_{1-4} alkyl- group in which the amino is optionally substituted by one or more C_{1-3} alkyl groups;

an optionally substituted non-aromatic C_{3-15} carbocyclic group;

a (C_{3-12} cycloalkyl) C_{1-3} alkyl- group;

a group $-(CH_2)_r(phenyl)_s$, in which r is 0,1, 2, 3 or 4, s is 1 when r is 0 otherwise s is 1 or 2

10 and the phenyl groups are optionally independently substituted by one, two or three groups represented by Z;

naphthyl;

anthracenyl;

a saturated 5 to 8 membered heterocyclic group containing one nitrogen and optionally

15 one of the following : oxygen, sulphur or an additional nitrogen wherein the heterocyclic group is optionally substituted by one or more C_{1-3} alkyl groups, hydroxy or benzyl ;

1-adamantylmethyl;

a group $-(CH_2)_t Het$ in which t is 0,1, 2, 3 or 4, and the alkylene chain is optionally substituted by one or more C_{1-3} alkyl groups and Het represents an aromatic heterocycle

20 optionally substituted by one, two or three groups selected from a C_{1-5} alkyl group, a C_{1-5} alkoxy group or halo;

or R^1 represents H and R^2 is as defined above;

or R^1 and R^2 together with the nitrogen atom to which they are attached represent a saturated 5 to 8 membered heterocyclic group containing one nitrogen and optionally one

25 of the following : oxygen, sulphur or an additional nitrogen; wherein the heterocyclic group is optionally substituted by one or more C_{1-3} alkyl groups, hydroxy or benzyl ;

X is CO or SO_2 ;

30 Y is absent or represents NH optionally substituted by a C_{1-3} alkyl group;

R^3 and R^4 independently represent phenyl, thienyl or pyridyl each of which is optionally substituted by one, two or three groups represented by Z;

5 Z represents a C_{1-3} alkyl group, a C_{1-3} alkoxy group, hydroxy, halo, trifluoromethyl, trifluoromethylthio, trifluoromethoxy, trifluoromethylsulphonyl, nitro, amino, mono or di C_{1-3} alkylamino, mono or di C_{1-3} alkylamido, C_{1-3} alkylsulphonyl, C_{1-3} alkoxycarbonyl, carboxy, cyano, carbamoyl, mono or di C_{1-3} alkyl carbamoyl, sulphamoyl and acetyl; and

10 R^5 is H, a C_{1-3} alkyl group, a C_{1-3} alkoxymethyl group, trifluoromethyl, a hydroxy C_{1-3} alkyl group, C_{1-3} alkoxycarbonyl, carboxy, cyano, carbamoyl, mono or di C_{1-3} alkylcarbamoyl, acetyl, or hydrazinocarbonyl of formula $-\text{CONHNR}^a\text{R}^b$ wherein R^a and R^b are as previously defined for R^1 and R^2 respectively;

15 with the proviso that when R^1 and R^2 together with the nitrogen atom to which they are attached represent 4-methylpiperazin-1-yl or R^1 represents H and R^2 represents methyl or 1-benzylpiperidin-4-yl; X is CO; Y is absent and R^5 is H; then R^3 and R^4 do not both represent 4-methoxyphenyl.


20 Further values of R^1 , R^2 , R^3 , R^4 and R^5 in compounds of formula I now follow. It will be understood that such values may be used where appropriate with any of the definitions, claims or embodiments defined hereinbefore or hereinafter.

In one group of compounds of formula I, R^1 represents H, R^2 represents cyclohexyl, X is
25 CO and Y is absent.

In a second group of compounds of formula I, R^1 and R^2 together with the nitrogen atom to which they are attached represent 1-piperidinyl.

30 In a third group of compounds of formula I, R^1 represents H and R^2 represents phenyl.

A fourth group of compounds of formula I is represented by formula Ia



5 and pharmaceutically acceptable salts, solvates and crystalline forms thereof, in which

R^2 represents cyclohexyl, 1-piperidinyl or phenyl ;

R^6 represents H, chloro, bromo, methyl or methoxy; and

when R^7 represents H, R^8 represents H or chloro; and

when R^7 represents chloro, R^8 represents H or chloro.

10

In a fifth group of compounds of formula I R^5 is H.

In a sixth group of compounds of formula I X is CO.

15 In a seventh group of compounds of formula I X is SO_2 .

In an eighth group of compounds of formula I Y is absent.

“Pharmaceutically acceptable salt”, where such salts are possible, includes both

20 pharmaceutically acceptable acid and base addition salts. A suitable pharmaceutically acceptable salt of a compound of Formula I is, for example, an acid-addition salt of a compound of Formula I which is sufficiently basic, for example an acid-addition salt with an inorganic or organic acid such as hydrochloric, hydrobromic, sulphuric, trifluoroacetic, citric or maleic acid; or, for example a salt of a compound of Formula I which is

25 sufficiently acidic, for example an alkali or alkaline earth metal salt such as a sodium,

calcium or magnesium salt, or an ammonium salt, or a salt with an organic base such as methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

5 Throughout the specification and the appended claims, a given chemical formula or name shall encompass all stereo and optical isomers and racemates thereof as well as mixtures in different proportions of the separate enantiomers, where such isomers and enantiomers exist, as well as pharmaceutically acceptable salts thereof and solvates thereof such as for instance hydrates. Isomers may be separated using conventional techniques, e.g.
10 chromatography or fractional crystallisation. The enantiomers may be isolated by separation of racemate for example by fractional crystallisation, resolution or HPLC. The diastereomers may be isolated by separation of isomer mixtures for instance by fractional crystallisation, HPLC or flash chromatography. Alternatively the stereoisomers may be made by chiral synthesis from chiral starting materials under conditions which will not
15 cause racemisation or epimerisation, or by derivatisation, with a chiral reagent. All stereoisomers are included within the scope of the invention. All tautomers, where possible, are included within the scope of the invention.

The following definitions shall apply throughout the specification and the appended
20 claims.

Unless otherwise stated or indicated, the term "alkyl" denotes either a straight or branched alkyl group. Examples of said alkyl include methyl, ethyl, n-propyl, isopropyl, n-butyl, iso-butyl, sec-butyl and t-butyl. Preferred alkyl groups are methyl, ethyl, propyl,
25 isopropyl and tertiary butyl.

Unless otherwise stated or indicated, the term "alkoxy" denotes a group O-alkyl, wherein alkyl is as defined above.

30 Unless otherwise stated or indicated, the term "halogen" shall mean fluorine, chlorine, bromine or iodine.

Specific compounds of the invention are one or more of the following:

- N*-(1-piperidinyl)-5,6-diphenyl-2-pyrazinecarboxamide;
N-(1-piperidinyl)-5,6-bis(4-bromophenyl)-2-pyrazinecarboxamide;
N-(1-piperidinyl)-5,6-bis(4-methylphenyl)-2-pyrazinecarboxamide;
5 *N*-(1-piperidinyl)-5,6-bis(4-methoxyphenyl)-2-pyrazinecarboxamide;
N-(1-piperidinyl)-5,6-bis(4-chlorophenyl)-2-pyrazinecarboxamide;
N-(1-piperidinyl)-5,6-bis(2-chlorophenyl)-2-pyrazinecarboxamide;
N-cyclohexyl-5,6-diphenyl-2-pyrazinecarboxamide;
N-cyclohexyl-5,6-bis(4-bromophenyl)-2-pyrazinecarboxamide;
10 *N*-cyclohexyl-5,6-bis(4-methylphenyl)-2-pyrazinecarboxamide;
N-cyclohexyl-5,6-bis(4-methoxyphenyl)-2-pyrazinecarboxamide;
N-cyclohexyl-5,6-bis(4-chlorophenyl)-2-pyrazinecarboxamide;
N-cyclohexyl-5,6-bis(2-chlorophenyl)-2-pyrazinecarboxamide;
N,5,6-triphenyl-2-pyrazinecarboxamide;
15 *N*-phenyl-5,6-bis(4-methylphenyl)-2-pyrazinecarboxamide;
N-phenyl-5,6-bis(4-methoxyphenyl)-2-pyrazinecarboxamide;
N-phenyl-5,6-bis(4-chlorophenyl)-2-pyrazinecarboxamide;
N-phenyl-5,6-bis(2-chlorophenyl)-2-pyrazinecarboxamide;
N-(1-piperidinyl)-5-(4-chlorophenyl)-6-(2,4-dichlorophenyl)-2-pyrazinecarboxamide; and
20 *N*-(1-piperidinyl)-6-(4-chlorophenyl)-5-(2,4-dichlorophenyl)-2-pyrazinecarboxamide;
and where applicable, optical isomers, tautomers, stereoisomers and racemates thereof as
well as pharmaceutically acceptable salts, solvates and crystalline forms thereof.


Methods of preparation

25

The compounds of the invention may be prepared as outlined below according to any of the following methods. However, the invention is not limited to these methods, the compounds may also be prepared as described for structurally related compounds in the prior art.


30

Compounds of formula I in which X is CO may be prepared by reacting a compound of formula II




II

in which R^3 , R^4 and R^5 are as previously defined with an amine of formula III




in an inert solvent, for example dichloromethane, in the presence of a coupling agent, for example a carbodiimide, eg 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide, and optionally in the presence of a catalyst, for example a basic catalyst, eg 4-dimethylamino-
pyridine, at a temperature in the range of -25°C to 150°C .

Compounds of formula I in which X is SO_2 may be prepared by reacting a compound of formula IV



IV

in which R^3 , R^4 and R^5 are as previously defined and A represents halo with an amine of formula IV



in an inert solvent, for example dichloromethane, and optionally in the presence of a catalyst, for example a basic catalyst, eg 4-dimethylamino-pyridine, at a temperature in the range of -25°C to 150°C .

Compounds of formulae II, III, IV and V may be prepared as described in the Examples and by other methods known to those skilled in the art. Certain compounds of formulae II, III, IV and V are novel and are claimed as a further aspect of the present invention as
5 useful intermediates. Specifically claimed are compounds of formula II in which R^3 , R^4 and R^5 are as previously defined with the exception of 5,6-diphenyl-2-pyrazinecarboxylic acid and 5,6-bis(4-methoxyphenyl)-2-pyrazinecarboxylic acid.

The compounds of the invention may be isolated from their reaction mixtures using
10 conventional techniques.

Persons skilled in the art will appreciate that, in order to obtain compounds of the invention in an alternative and in some occasions, more convenient manner, the individual process steps mentioned hereinbefore may be performed in different order, and/or the individual
15 reactions may be performed at different stage in the overall route (i.e. chemical transformations may be performed upon different intermediates to those associated hereinbefore with a particular reaction).

The expression "inert solvent" refers to a solvent which does not react with the starting
20 materials, reagents, intermediates or products in a manner which adversely affects the yield of the desired product.

Pharmaceutical preparations

The compounds of the invention will normally be administered via the oral, parenteral, intravenous, intramuscular, subcutaneous or in other injectable ways, buccal, rectal,
25 vaginal, transdermal and/or nasal route and/or via inhalation, in the form of pharmaceutical preparations comprising the active ingredient or a pharmaceutically acceptable addition salt, in a pharmaceutically acceptable dosage form. Depending upon the disorder and patient to be treated and the route of administration, the compositions may be administered at varying doses.

Suitable daily doses of the compounds of the invention in the therapeutic treatment of humans are about 0.001-10 mg/kg body weight, preferably 0.01-1 mg/kg body weight.

Oral formulations are preferred particularly tablets or capsules which may be formulated by methods known to those skilled in the art to provide doses of the active compound in the range of 0.5mg to 500mg for example 1 mg, 3 mg, 5 mg, 10 mg, 25mg, 50mg, 100mg and 250mg.

A compound of the invention may also be combined with other anti-obesity agents such as Orlistat or a monoamine reuptake inhibitor, for example Sibutramine. Furthermore, a compound of the invention may also be combined with therapeutic agents that are useful in the treatment of disorders or conditions associated with obesity (such as type II diabetes, metabolic syndrome, dyslipidemia, impaired glucose tolerance, hypertension, coronary heart disease, non-alcoholic steatorrheic hepatitis, osteoarthritis and some cancers) and psychiatric and neurological conditions.

According to a further aspect of the invention there is also provided a pharmaceutical formulation including any of the compounds of the invention, or pharmaceutically acceptable derivatives thereof, in admixture with pharmaceutically acceptable adjuvants, diluents and/or carriers.

Pharmacological properties

The compounds of formula (I) are useful for the treatment of obesity, psychiatric disorders such as psychotic disorders, schizophrenia, bipolar disorders, anxiety, anxio-depressive disorders, depression, cognitive disorders, memory disorders, obsessive-compulsive disorders, anorexia, bulimia, attention disorders like ADHD, epilepsy, and related conditions, and neurological disorders such as dementia, neurological disorders(e.g. Multiple Sclerosis), Raynaud's syndrome, Parkinson's disease, Huntington's chorea and Alzheimer's disease. The compounds are also potentially useful for the treatment of immune, cardiovascular, reproductive and endocrine disorders, septic shock and diseases related to the respiratory and gastrointestinal systems (e.g. diarrhea). The compounds are


also potentially useful as agents in treatment of extended abuse, addiction and/or relapse indications, e.g. treating drug (nicotine, ethanol, cocaine, opiates, etc) dependence and/or treating drug (nicotine, ethanol, cocaine, opiates, etc) withdrawal symptoms. The compounds may also eliminate the increase in weight which normally accompanies the
5 cessation of smoking.

In another aspect the present invention provides a compound of formula I as previously defined for use as a medicament.

10 In a further aspect the present invention provides the use of a compound of formula I (including the compounds of the proviso) in the preparation of a medicament for the treatment or prophylaxis of obesity, psychiatric disorders such as psychotic disorders, schizophrenia, bipolar disorders, anxiety, anxio-depressive disorders, depression, cognitive disorders, memory disorders, obsessive-compulsive disorders, anorexia, bulimia, attention
15 disorders like ADHD, epilepsy, and related conditions, neurological disorders such as dementia, neurological disorders (e.g. Multiple Sclerosis), Parkinson's Disease, Huntington's Chorea and Alzheimer's Disease, immune, cardiovascular, reproductive and endocrine disorders, septic shock, diseases related to the respiratory and gastrointestinal systems (e.g. diarrhea), and extended abuse, addiction and/or relapse indications, e.g.
20 treating drug (nicotine, ethanol, cocaine, opiates, etc) dependence and/or treating drug (nicotine, ethanol, cocaine, opiates, etc) withdrawal symptoms.

In a still further aspect the present invention provides a method of treating obesity, psychiatric disorders such as psychotic disorders such as schizophrenia and bipolar
25 disorders, anxiety, anxio-depressive disorders, depression, cognitive disorders, memory disorders, obsessive-compulsive disorders, anorexia, bulimia, attention disorders like ADHD, epilepsy, and related conditions, neurological disorders such as dementia, neurological disorders (e.g. Multiple Sclerosis), Parkinson's Disease, Huntington's Chorea and Alzheimer's Disease, immune, cardiovascular, reproductive and endocrine disorders,
30 septic shock, diseases related to the respiratory and gastrointestinal systems (e.g. diarrhea), and extended abuse, addiction and/or relapse indications, e.g. treating drug (nicotine, ethanol, cocaine, opiates, etc) dependence and/or treating drug (nicotine, ethanol, cocaine,

opiates, etc) withdrawal symptoms comprising administering a pharmacologically effective amount of a compound of formula I



I

and pharmaceutically acceptable salts, prodrugs, solvates and crystalline forms thereof, in which

R^1 and R^2 independently represent :

a C_{1-6} alkyl group;

an (amino) C_{1-4} alkyl- group in which the amino is optionally substituted by one or more

C_{1-3} alkyl groups;

an optionally substituted non-aromatic C_{3-15} carbocyclic group;

a (C_{3-12} cycloalkyl) C_{1-3} alkyl- group;

a group $-(CH_2)_r(phenyl)_s$ in which r is 0,1, 2, 3 or 4, s is 1 when r is 0 otherwise s is 1 or 2

and the phenyl groups are optionally independently substituted by one, two or three groups

represented by Z;

naphthyl;

anthracenyl;

a saturated 5 to 8 membered heterocyclic group containing one nitrogen and optionally one of the following : oxygen, sulphur or an additional nitrogen wherein the heterocyclic

group is optionally substituted by one or more C_{1-3} alkyl groups, hydroxy or benzyl ;

1-adamantylmethyl;

a group $-(CH_2)_t$ Het in which t is 0,1, 2, 3 or 4, and the alkylene chain is optionally substituted by one or more C_{1-3} alkyl groups and Het represents an aromatic heterocycle optionally substituted by one, two or three groups selected from a C_{1-5} alkyl group, a

C_{1-5} alkoxy group or halo;

or R^1 represents H and R^2 is as defined above;

or R^1 and R^2 together with the nitrogen atom to which they are attached represent a saturated 5 to 8 membered heterocyclic group containing one nitrogen and optionally one of the following : oxygen, sulphur or an additional nitrogen; wherein the heterocyclic group is optionally substituted by one or more C_{1-3} alkyl groups, hydroxy or benzyl ;

X is CO or SO₂ ;

Y is absent or represents NH optionally substituted by a C_{1-3} alkyl group;

R^3 and R^4 independently represent phenyl, thienyl or pyridyl each of which is optionally substituted by one , two or three groups represented by Z;

Z represents a C_{1-3} alkyl group, a C_{1-3} alkoxy group, hydroxy, halo, trifluoromethyl, trifluoromethylthio, trifluoromethoxy, trifluoromethylsulphonyl, nitro, amino, mono or di C_{1-3} alkylamino, mono or di C_{1-3} alkylamido, C_{1-3} alkylsulphonyl, C_{1-3} alkoxycarbonyl, carboxy, cyano, carbamoyl, mono or di C_{1-3} alkyl carbamoyl, sulphamoyl and acetyl; and

R^5 is H, a C_{1-3} alkyl group, a C_{1-3} alkoxymethyl group, trifluoromethyl, a hydroxy C_{1-3} alkyl group, C_{1-3} alkoxycarbonyl, carboxy, cyano, carbamoyl, mono or di C_{1-3} alkylcarbamoyl, acetyl, or hydrazinocarbonyl of formula $-\text{CONHNR}^a\text{R}^b$ wherein R^a and R^b are as previously defined for R^1 and R^2 respectively;

to a patient in need thereof.

The compounds of the present invention are particularly suitable for the treatment of obesity, e.g. by reduction of appetite and body weight, maintenance of weight reduction and prevention of rebound.

ExamplesAbbreviations

DCM - dichloromethane

5 DMF - dimethylformamide

DMAP - 4-dimethylaminopyridine

EDC - 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide

TEA - triethylamine

TFA - trifluoroacetic acid

10 DMSO - dimethyl sulfoxide

DEA - Diethylamine

PCC - Pyridinium chlorochromate

DCM - Dichloromethane

15	t	triplet
	s	singlet
	d	doublet
	q	quartet
	qvint	quintet
20	m	multiplet
	br	broad
	bs	broad singlet
	dm	doublet of multiplet
	bt	broad triplet
25	dd	doublet of doublet

General Experimental Procedures

Mass spectra were recorded on either a Micromass ZQ single quadrupole or a Micromass LCZ single quadrupole mass spectrometer both equipped with a pneumatically assisted electrospray interface (LC-MS). ¹H NMR measurements were performed on either a
30 Varian Mercury 300 or a Varian Inova 500, operating at ¹H frequencies of 300 and 500 MHz respectively. Chemical shifts are given in ppm with CDCl₃ as internal standard.

Purification was performed on a semipreparative HPLC with a mass triggered fraction collector, Shimadzu QP 8000 single quadrupole mass spectrometer equipped with 19 x 100 mm C8 column. The mobile phase used was, if nothing else is stated, acetonitrile and buffer (0.1 M NH₄Ac:acetonitrile 95:5).

For isolation of isomers, a Kromasil CN E9344 (250 x 20 mm i.d.) column was used. Heptane:ethyl acetate:DEA 95:5:0.1 was used as mobile phase (1 ml/min). Fraction collection was guided using a UV-detector (330 nm).

10 Synthesis of intermediates

The following intermediates were not commercially available and therefore prepared as described in Preparation A, (Chem. Ber., 100, 1967, p. 555).

Preparation A

15 (a) 5,6-diphenyl-pyrazine-2-carboxylic acid

The monohydrochloride of 2,3-diaminopropionic acid (500 mg, 3.56 mmol) and benzil (890 mg, 4.23 mmol) were added to a solution of sodium hydroxide (677 mg, 16.93 mmol) in methanol (10 ml). An extra portion of methanol was added (5 ml) and the reaction mixture was refluxed for 20 minutes. The mixture was cooled to 25°C and air was bubbled through for 30 minutes. Hydrochloric acid (aq, 2 M) was added until the reaction mixture reached pH 2. The solution was extracted with diethyl ether. The combined diethyl ether phases were dried (MgSO₄), filtrated and evaporated under reduced pressure to give the crude product. MS *m/z* 277 (M+H)⁺. The crude product was used in steps described below without further purification.

25

(b) 5,6-Bis-(4-bromophenyl)-pyrazine-2-carboxylic acid

The title compound was prepared essentially as described in Preparation A step (a), using monohydrochloride of 2,3-diaminopropionic acid (600 mg, 4.26 mmol) and 4,4'-dibromobenzil (1.745 g, 4.26 mmol, 90 %) as starting materials. The reaction mixture was refluxed for 2 hours and air was bubbled through for 1 hour. Hydrochloric acid (aq, 2 M)

30

was added until pH 2. The mixture was evaporated under reduced pressure and the residue was dissolved in water. The solution was extracted with diethyl ether, the combined diethyl ether phases were dried (MgSO_4), filtered and evaporated under reduced pressure. The crude product (500 mg, 27%) was used in steps described below without further purification. MS m/z 435, 437, 439 ($\text{M}+\text{H}$)⁺.

(c) 5,6-Di-*p*-tolyl-pyrazine-2-carboxylic acid

The title compound was prepared as described in Preparation A step (a) using 4,4'-dimethylbenzil (848 mg, 3.56 mmol). The reaction mixture was however refluxed for 1 hour and air was bubbled through the reaction mixture for about 7 hours. The mixture was evaporated and the residue was dissolved in water. Hydrochloric acid (aq, 2 M) was added until pH 2 was reached. The solution was extracted with diethyl ether. The combined diethyl ether phases were dried (MgSO_4), filtered and evaporated under reduced pressure. The crude product (918 mg, 85%) was used in steps described below without further purification. MS m/z 305 ($\text{M}+\text{H}$)⁺.

(d) 5,6-Bis-(4-methoxyphenyl)pyrazine-2-carboxylic acid

The title compound was prepared as described in Preparation A step (c) using 4,4'-dimethoxybenzil (961 mg, 3.56 mmol) as starting material. The reaction mixture was refluxed over night and air was bubbled through the mixture for 8 hours. The crude product (435 mg, 36%) was used in steps described below without further purification. MS m/z 335 ($\text{M}+\text{H}$)⁺.

(e) 5,6-Bis-(4-chlorophenyl) pyrazine-2-carboxylic acid

The title compound was prepared as described in Preparation A step (c) using 4,4'-dichlorobenzil (993 mg, 3.56 mmol). Reflux for 1 hour gave directly the crude product (923 mg, 75%) that was used in steps described below without further purification. MS m/z 343, 345, 347 ($\text{M}-\text{H}$)⁻.

(f) 5,6-Bis-(2-chlorophenyl)pyrazine-2-carboxylic acid

The title compound was prepared as described in Preparation A step (c) using 2,2'-dichlorobenzil (993 mg, 3.56 mmol). The crude product (895 mg, 73%) was used in steps described below without further purification. MS m/z 343, 345, 347 (M-H).

5 (h) 1-(4-Chlorophenyl)-2-(2,4-dichlorophenyl)ethane-1,2-dione
2-(4-Chlorophenyl)-1-(2,4-dichlorophenyl)ethanone (2.7 g, 9.01 mmol) was dissolved in 1,2-dichloroethane (25 ml) and freshly made PCC (3.89 g, 18.02 mmol), pyridine (1.43 g, 18.02 mmol) and molecular sieves were added. The reaction mixture was refluxed under inert atmosphere overnight. The solution was cooled to 25 °C, filtered through Silica and
10 then solvent was evaporated under reduced pressure. The crude product (1.9 g, 66%) was used directly in the next step. ¹H NMR (500 MHz) δ 7.97 (d, 2H), 7.84 (d, 1H), 7.52 (d, 2H), 7.46 (s, 1H), 7.44 (d, 1H).

(i) 5-(4-Chlorophenyl)-6-(2,4-dichlorophenyl)pyrazine-2-carboxylic acid and
15 6-(4-chlorophenyl)-5-(2,4-dichlorophenyl)pyrazine-2-carboxylic acid
The title compounds were prepared as described in Preparation A step (a), using 1-(4-chlorophenyl)-2-(2,4-dichlorophenyl)ethane-1,2-dione (1.85 g; 5.90 mmol) from Preparation A step (g) and the monochloride of 2,3-diaminopropionic acid (0.61 g, 5.90 mmol) as starting materials. The mixture was refluxed for 30 minutes and then directly
20 worked-up. The crude product was allowed to stand over night to aromatise. Flash chromatography (SiO₂, DCM:methanol 10:1, 1% Acetic acid) gave the isomer mixture (0.2 g, 10%). MS m/z 377, 379, 381 (M-H).

Examples of the Invention

25

Example 1

N-(1-piperidinyl)- 5,6-diphenyl-2-pyrazinecarboxamide

5,6-Diphenyl-pyrazine-2-carboxylic acid (500 mg, 1.81 mmol) from Preparation A, step (a), was dissolved in DCM (4 ml) and DMF (150 μ l). DMAP (22 mg, 0.18 mmol) and 1-aminopiperidine (218 mg, 2.17 mmol) were added and the solution was cooled to 0 °C. A
30 slurry of EDC (1.99 mmol, in 2mL DCM and 100 μ l DMF) was added dropwise. The

reaction mixture was stirred at 25 °C. After 17 hours additional 1-aminopiperidine (40 mg, 0.40 mmol) and EDC (76 mg, 0.40 mmol) was added, and the mixture was stirred for an additional 3 hours. The crude was diluted with DCM (5ml) and washed with a saturated solution of NaHCO₃. The organic phase was dried (MgSO₄), filtered and evaporated. Flash chromatography (SiO₂, ethyl acetate:hexane 2:1) gave the subtitle compound (160 mg, 25%) as a white solid.

¹H NMR (300 MHz) δ 9.41 (s, 1H), 8.52 (s, 1H), 7.50-7.29 (m, 10H), 2.94 (t, 4H), 1.81 (m, 4H), 1.50 (m, 2H).

MS *m/z* 359 (M+H)⁺.

Example 2

N-(1-piperidinyl)- 5,6-bis(4-bromophenyl)-2-pyrazinecarboxamide

To 5,6-Bis-(4-bromophenyl)-pyrazine-2-carboxylic acid (108 mg, 0.25 mmol) from Preparation A, step (b), DMAP (0.025 mmol, in 500 µl DCM), 1-aminopiperidine (0.25 mmol, in 1100 µl DCM), EDC (0.27 mmol, in 1100 µl DCM and cooled to 8 °C) were added. The reaction mixture was stirred at 25 °C for 20 h, then washed with saturated NaHCO₃ solution, dried (MgSO₄), filtered and evaporated. Semipreparatory HPLC (0.01% TEA in the buffered phase) gave the subtitle compound (6.7 mg, 5.4%).

¹H NMR (300 MHz) δ 9.41 (s, 1H), 8.48 (s, 1H), 7.54 (d, 2H), 7.51 (d, 2H), 7.36 (d, 2H), 7.34 (d, 2H), 2.94 (t, 4H), 1.81 (m, 4H), 1.55-1.45 (m, 2H).

MS *m/z* 515, 517, 519 (M+H)⁺.

Example 3

N-(1-piperidinyl)- 5,6-bis(4-methylphenyl)- 2-pyrazinecarboxamide

5,6-Di-*p*-tolyl-pyrazine-2-carboxylic acid (76 mg, 0.25 mmol) from Preparation A, step (c), was used as described in Example 2 to give the title compound (27 mg, 28%).

¹H NMR (300 MHz) δ 9.35 (s, 1H), 8.57 (s, 1H), 7.38 (d, 4H), 7.18 (d, 2H), 7.13 (d, 2H), 2.92 (t, 4H), 2.40 (s, 3H), 2.37 (s, 3H), 1.86-1.75 (m, 4H), 1.54-1.44 (m, 2H).

MS *m/z* 387 (M+H)⁺.

Example 4*N*-(1-piperidinyl)-5,6-bis(4-methoxyphenyl)-2-pyrazinecarboxamide

5,6-Bis-(4-methoxyphenyl)-pyrazine-2-carboxylic acid (84 mg, 0.25 mmol) from Preparation A, step (d), was used as described Example 2 to give the title compound (20 mg, 19%).

¹H NMR (300 MHz) δ 9.31 (s, 1H), 8.57 (s, 1H), 7.46 (d, 2H), 7.44 (d, 2H), 6.90 (d, 2H), 6.86 (d, 2H), 3.86 (s, 3H), 3.84 (s, 3H), 2.93 (t, 4H), 1.80 (m, 4H), 1.54-1.45 (m, 2H).

MS *m/z* 419 (M+H)⁺.

Example 5*N*-(1-piperidinyl)-5,6-bis(4-chlorophenyl)-2-pyrazinecarboxamide

5,6-Bis-(4-chlorophenyl)-pyrazine-2-carboxylic acid (86 mg, 0.25 mmol) from Preparation A, step (e), was used as described in Example 2 to give the subtitle compound (16 mg, 15%).

¹H NMR (300 MHz) δ 9.40 (s, 1H), 8.49 (s, 1H), 7.45-7.31 (m, 8H), 2.94 (t, 4H), 1.80 (m, 4H), 1.54-1.45 (m, 2H).

MS *m/z* 427, 429, 431 (M+H)⁺.

Example 6*N*-(1-piperidinyl)-5,6-bis(2-chlorophenyl)-2-pyrazinecarboxamide

5,6-Bis-(2-chlorophenyl)-pyrazine-2-carboxylic acid (86 mg, 0.25 mmol) from Preparation A, step (f), was used as described in Example 2 to give the subtitle compound (6 mg, 6%).

¹H NMR (300 MHz) δ 9.52 (s, 1H), 8.52 (s, 1H), 7.44-7.17 (d, 8H), 2.94-2.88 (t, 4H), 1.85-1.70 (m, 4H), 1.52-1.44 (m, 2H).

MS *m/z* 427, 429, 431 (M+H)⁺.

Example 7*N*-cyclohexyl-5,6-diphenyl-2-pyrazinecarboxamide

5,6-diphenyl-pyrazine-2-carboxylic acid (70 mg, 0.25 mmol) from Preparation A, step (a), was reacted essentially as described in Example 2 but with cyclohexylamine (0.25 mmol,

in 1 ml DCM), DMAP (0.025 mmol, in 0.5 ml DCM), EDC (0.28 mmol, in 1 ml DCM, and cooled to 8 °C) and DMF (100 µl). Semipreparatory HPLC (0.15% TFA/water:acetonitrile 95:5 instead of the buffer phase) gave the title compound (7 mg, 8%) after washing with Na₂CO₃ solution.

5 ¹H NMR (300 MHz) δ 9.41 (s, 1H), 7.78 (d, 1H), 7.49-7.28 (m, 10H), 4.12-3.97 (m, 1H), 2.13-1.23 (m, 10H).

MS *m/z* 358 (M+H)⁺.

Example 8

10 *N*-cyclohexyl-5,6-bis(4-bromophenyl)-2-pyrazinecarboxamide

5,6-Bis-(4-bromophenyl)-pyrazine-2-carboxylic acid (109 mg, 0.25 mmol) from Preparation A, step (b), was used as described in Example 7. Semipreparatory HPLC (0.15 % TFA/water:acetonitrile 95:5 instead of the buffer phase) gave the title compound (7 mg, 8%) after washing with Na₂CO₃ solution.

15 ¹H NMR (300 MHz) δ 9.41 (s, 1H), 7.68 (s, 1H), 7.54 (d, 2H), 7.50 (d, 2H), 7.36 (d, 2H), 7.34 (d, 2H), 4.11-3.96(m, 1H), 2.12-1.20 (m, 10H).

MS *m/z* 514, 516, 518 (M+H)⁺.

Example 9

20 *N*-cyclohexyl-5,6-bis(4-methylphenyl)-2-pyrazinecarboxamide

5,6-Di-*p*-tolyl-pyrazine-2-carboxylic acid (76 mg, 0.25 mmol) from Preparation A, step (c), was used as described in Example 7. Semipreparatory HPLC (0.01% TEA in the buffer phase) gave the subtitle compound (4 mg, 4%).

¹H NMR (300 MHz) δ 9.36 (s, 1H), 7.77 (d, 1H), 7.39 (d, 4H), 7.18 (d, 2H), 7.13 (d, 2H),
25 4.10-3.96 (m, 1H), 2.40 (s, 3H), 2.37 (s, 3H), 2.09-1.20 (m, 10H).

MS *m/z* 386 (M+H)⁺.

Example 10N-cyclohexyl-5,6-bis(4-methoxyphenyl)-2-pyrazinecarboxamide

5,6-Bis-(4-methoxyphenyl)-pyrazine-2-carboxylic acid (76 mg, 0.25 mmol) from

Preparation A, step (d), was used essentially as described in Example 7 but the reaction

5 mixture was first stirred overnight, then more cyclohexylamine (25 mg, 0.25 mmol) was added and the mixture was stirred for an additional two days prior to workup.

Semipreparatory HPLC (0.15 % TFA in the buffered phase) gave the title compound (12 mg, 11%).

¹H NMR (300 MHz) δ 9.32 (s, 1H), 7.76 (d, 1H), 7.47 (d, 2H), 7.45 (d, 2H), 6.90 (d, 2H),
10 6.86 (d, 2H), 4.10-3.96 (m, 1H), 3.86 (s, 3H), 3.84 (s, 3H), 2.09-1.17 (m, 10H).

MS *m/z* 418 (M+H)⁺.

Example 11N-cyclohexyl-5,6-bis(4-chlorophenyl)-2-pyrazinecarboxamide

15 5,6-Bis-(4-chlorophenyl)pyrazine-2-carboxylic acid (86 mg, 0.25 mmol) from Preparation A, step (e), was used as described in Example 10 to give the title compound (7 mg, 8%) after washing with Na₂CO₃ solution.

¹H NMR (300 MHz) δ 9.41 (s, 1H), 7.69 (s, 1H), 7.47-7.30 (m, 8H), 4.10-3.97 (m, 1H),
2.10-1.18 (m, 10H).

20 MS *m/z* 426, 428, 430 (M+H)⁺.

Example 12N-cyclohexyl-5,6-bis(2-chlorophenyl)-2-pyrazinecarboxamide

5,6-Bis-(2-chlorophenyl)-pyrazine-2-carboxylic acid (86 mg, 0.25 mmol) from preparation

25 A step (f) was used as described in Example 10, to give the title compound (14 mg, 13%).

¹H NMR (300 MHz) δ 9.51 (s, 1H), 7.74 (s, 1H), 7.41-7.18 (m, 8H), 4.10-3.97 (m, 1H),
2.07-1.14 (m, 10H).

MS *m/z* 426, 428, 430 (M+H)⁺.

Example 13N,5,6-triphenyl-2-pyrazinecarboxamide

To 5,6-Diphenyl-pyrazine-2-carboxylic acid (70 mg, 0.25 mmol) from Preparation A, step (a), DMAP (0.025 mmol, in 0.5 ml DCM), aniline (0.25 mmol, in 1 ml DCM), EDC (0.28 mmol, in 1 ml DCM, cooled to 8 °C) and DMF (100 µl) were added. The reaction mixture was stirred at 25 °C over night, then worked up as described in Example 2.

Semipreparatory HPLC (0.15 % TFA/water:acetonitrile 95:5 instead of the buffer phase) gave the title compound (27 mg, 30%) after washing with Na₂CO₃ solution.

¹H NMR (300 MHz) δ 9.75 (s, 1H), 9.52 (d, 1H), 7.80 (d, 2H), 7.55-7.32 (m, 12H), 7.20 (t, 1H).

MS *m/z* 352 (M+H)⁺.

Example 14N-phenyl-5,6-bis(4-methylphenyl)-2-pyrazinecarboxamide

5,6-Di-*p*-tolyl-pyrazine-2-carboxylic acid (77 mg, 0.25 mmol) from Preparation A, step (c), was used as described in Example 13 to give the subtitle compound (28 mg, 29%).

¹H NMR (500 MHz) δ 9.78 (s, 1H), 9.49 (s, 1H), 7.81 (d, 2H), 7.47-7.43 (m, 6H), 7.25-7.17 (m, 5H), 2.45 (s, 3H), 2.41 (s, 3H).

MS *m/z* 380 (M+H)⁺.

Example 15N-phenyl-5,6-bis(4-methoxyphenyl)-2-pyrazinecarboxamide

5,6-Bis-(4-methoxyphenyl)-pyrazine-2-carboxylic acid (85 mg, 0.25 mmol) from Preparation A step (d), was used as described in Example 13, to give the title compound (33 mg, 32%).

¹H NMR (300 MHz) δ 9.74 (s, 1H), 9.42 (s, 1H), 7.79 (d, 2H), 7.50 (d, 4H), 7.42 (t, 2H), 7.19 (t, 1H), 6.94 (d, 2H), 6.89 (d, 2H), 3.88 (s, 3H), 3.85 (s, 3H).

MS *m/z* 412 (M+H)⁺.

Example 16N-phenyl-5,6-bis(4-chlorophenyl)-2-pyrazinecarboxamide

5,6-Bis-(4-chlorophenyl)-pyrazine-2-carboxylic acid (87 mg, 0.25 mmol) from Preparation A, step (e), was used as described in Example 13, to give the subtitle compound (6 mg, 6%).

¹H NMR (300 MHz) δ 9.66 (s, 1H), 9.52 (s, 1H), 7.79 (d, 2H), 7.48-7.35 (m, 10H), 7.21 (t, 1H).

MS *m/z* 420, 422, 424 (M+H)⁺.

Example 17N-phenyl-5,6-bis(2-chlorophenyl)-2-pyrazinecarboxamide

5,6-Bis-(2-chloro-phenyl)-pyrazine-2-carboxylic acid (87 mg, 0.25 mmol) from Preparation A, step (f), was treated as described in Example 13, to give the title compound (27 mg, 25%).

¹H NMR (500 MHz) δ 9.73 (s, 1H), 9.66 (s, 1H), 7.81(d, 2H), 7.46-7.22 (m, 11H).

MS *m/z* 420, 422, 424 (M+H)⁺.

Example 185-(4-Chlorophenyl)-6-(2,4-dichlorophenyl)pyrazine-2-carboxylic acid piperidin-1-ylamide

20 and 6-(4-chlorophenyl)-5-(2,4-dichlorophenyl)pyrazine-2-carboxylic acid piperidin-1-ylamide

The mixture of 5-(4-chlorophenyl)-6-(2,4-dichlorophenyl)pyrazine-2-carboxylic acid and 6-(4-chlorophenyl)-5-(2,4-dichlorophenyl)pyrazine-2-carboxylic acid (78 mg, 0.205 mmol) from Preparation A step (i) and thionyl chloride (147 mg, 1.23 mmol) were refluxed in toluene (2ml) for 3 hours. The solvent and reagents were evaporated under reduced pressure and the intermediates were dissolved in DCM (1 ml). TEA (42 mg, 0.41 mmol) and 1-aminopiperidine (21 mg, 0.205 mmol) were dissolved in DCM (1ml) and added. The reaction mixture was stirred at 25 °C overnight and then evaporated under reduced pressure. Flash chromatography (SiO₂, heptane:ethyl acetate 1:1) gave a mixture of the title compounds (45 mg, 47%, ratio of isomers 0.5:1). ¹H NMR (300 MHz) δ 9.46 (s, 1H), 8.39 (s, 1H), 7.47-7.28 (m, 7H), 3.02-2.84 (m, 4H), 1.89-1.73 (m, 4H), 1.57-1.41 (m, 2H) and

9.42 (s, 1H), 8.51 (s, 1H), 7.47-7.28 (m, 7H), 3.02-2.84 (m, 4H), 1.89-1.73 (m, 4H), 1.57-1.41 (m, 2H).

Example 18 (a)

5 N-(1-piperidiny)- 5-(4-chlorophenyl)-6-(2,4-dichlorophenyl)-2-pyrazinecarboxamide

The title compound was isolated from the mixture prepared in Example 18 (35 mg) by preparative chromatography (9 mg, 26%). ¹H NMR (300 MHz) δ 9.46 (s, 1H), 8.38 (s, 1H), 7.46-7.24 (m, 7H), 2.89 (t, 4H), 1.78 (p, 4H), 1.52-1.40 (m, 2H).

10 Example 18 (b)

N-(1-piperidiny)- 6-(4-chlorophenyl)-5-(2,4-dichlorophenyl)-2-pyrazinecarboxamide

The title compound was isolated from the mixture prepared in Example 18 (35 mg) by preparative chromatography (11 mg, 31%). ¹H NMR (300 MHz) δ 9.42 (s, 1H), 8.50 (s, 1H), 7.39-7.30 (m, 7H), 2.93 (t, 4H), 1.80 (p, 4H), 1.54-1.43 (m, 2H).

15

Pharmacological Activity

Compounds of the present invention are active against the receptor product of the CB1 gene. The affinity of the compounds of the invention for central cannabinoid receptors is demonstrable in methods described in Devane et al , Molecular Pharmacology, 1988, 20 34,605 or those described in WO01/70700 or EP 656354. Alternatively the assay may be performed as follows.

10μg of membranes prepared from cells stably transfected with the CB1 gene were suspended in 200μl of 100mM NaCl, 5mM MgCl₂, 1mM EDTA, 50mM HEPES (pH 7.4), 1mM DTT, 0.1% BSA and 100μM GDP. To this was added an EC80 concentration of 25 agonist (CP55940), the required concentration of test compound and 0.1μCi [³⁵S]-GTPγS. The reaction was allowed to proceed at 30°C for 45 min. Samples were then transferred on to GF/B filters using a cell harvester and washed with wash buffer (50mM Tris (pH 7.4), 5mM MgCl₂, 50mM NaCl). Filters were then covered with scintillant and counted for the amount of [³⁵S]-GTPγS retained by the filter.

30 Activity is measured in the absence of all ligands (minimum activity) or in the presence of an EC80 concentration of CP55940 (maximum activity). These activities are set as 0% and 100% activity respectively. At various concentrations of novel ligand, activity is


calculated as a percentage of the maximum activity and plotted. The data are fitted using the equation $y=A+(B-A)/(1+((C/x)^D))$ and the IC₅₀ value determined as the concentration required to give half maximal inhibition of GTPγS binding under the conditions used.

5

The compounds of the present invention are active at the CB1 receptor (IC₅₀ <1 micromolar). Most preferred compounds have IC₅₀ <200 nanomolar.

Claims

1. A compound of formula (I)



and pharmaceutically acceptable salts, prodrugs, solvates and crystalline forms thereof, in which

R¹ and R² independently represent :

a C₁₋₆alkyl group;

an (amino)C₁₋₄alkyl- group in which the amino is optionally substituted by one or more C₁₋₃alkyl groups;

an optionally substituted non-aromatic C₃₋₁₅carbocyclic group;

a (C₃₋₁₂cycloalkyl)C₁₋₃alkyl- group;

a group -(CH₂)_r(phenyl)_s in which r is 0,1, 2, 3 or 4, s is 1 when r is 0 otherwise s is 1 or 2

and the phenyl groups are optionally independently substituted by one, two or three groups represented by Z;

naphthyl;

anthracenyl;

a saturated 5 to 8 membered heterocyclic group containing one nitrogen and optionally

one of the following : oxygen, sulphur or an additional nitrogen wherein the heterocyclic group is optionally substituted by one or more C₁₋₃alkyl groups, hydroxy or benzyl ;

1-adamantylmethyl;

a group -(CH₂)_t Het in which t is 0,1, 2, 3 or 4, and the alkylene chain is optionally substituted by one or more C₁₋₃alkyl groups and Het represents an aromatic heterocycle

optionally substituted by one, two or three groups selected from a C₁₋₅alkyl group, a C₁₋₅alkoxy group or halo;

or R¹ represents H and R² is as defined above;

or R^1 and R^2 together with the nitrogen atom to which they are attached represent a saturated 5 to 8 membered heterocyclic group containing one nitrogen and optionally one of the following : oxygen, sulphur or an additional nitrogen; wherein the heterocyclic group is optionally substituted by one or more C_{1-3} alkyl groups, hydroxy or benzyl ;

X is CO or SO_2 ;

Y is absent or represents NH optionally substituted by a C_{1-3} alkyl group;

R^3 and R^4 independently represent phenyl, thienyl or pyridyl each of which is optionally substituted by one , two or three groups represented by Z;

Z represents a C_{1-3} alkyl group, a C_{1-3} alkoxy group, hydroxy, halo, trifluoromethyl, trifluoromethylthio, trifluoromethoxy, trifluoromethylsulphonyl, nitro, amino, mono or di C_{1-3} alkylamino, mono or di C_{1-3} alkylamido, C_{1-3} alkylsulphonyl, C_{1-3} alkoxycarbonyl, carboxy, cyano, carbamoyl, mono or di C_{1-3} alkyl carbamoyl, sulphamoyl and acetyl; and

R^5 is H, a C_{1-3} alkyl group, a C_{1-3} alkoxymethyl group, trifluoromethyl, a hydroxy C_{1-3} alkyl group, C_{1-3} alkoxycarbonyl, carboxy, cyano, carbamoyl, mono or di C_{1-3} alkylcarbamoyl, acetyl, or hydrazinocarbonyl of formula $-CONHNR^aR^b$ wherein R^a and R^b are as previously defined for R^1 and R^2 respectively and;

with the proviso that when R^1 and R^2 together with the nitrogen atom to which they are attached represent 4-methylpiperazin-1-yl or R^1 represents H and R^2 represents methyl or 1-benzylpiperidin-4-yl; X is CO; Y is absent and R^5 is H; then R^3 and R^4 do not both represent 4-methoxyphenyl.


2. A compound according to claim 1 in which R^1 represents H, R^2 represents cyclohexyl, X is CO and Y is absent.

3. A compound according to claim 1 in which R^1 and R^2 together with the nitrogen atom to which they are attached represent 1-piperidinyl.

4. A compound according to claim 1 in which R^1 represents H and R^2 represents phenyl.

5

5. A compound according to claim 1 as represented by formula Ia



and pharmaceutically acceptable salts, solvates and crystalline forms thereof, in which

10 R^2 represents cyclohexyl, 1-piperidinyl or phenyl ;

R^6 represents H, chloro, bromo, methyl or methoxy; and

when R^7 represents H, R^8 represents H or chloro; and

when R^7 represents chloro, R^8 represents H or chloro.

15 6. A compound according to any one of claims 1 to 4 in which R^5 is H.

7. A compound according to any one of claims 1 to 4 in which X is CO.

8. A compound according to any one of claims 1 to 4 in which X is SO₂.

20


9. A compound according to any one of claims 1 to 4 in which Y is absent.

10. A compound selected from:

N-(1-piperidinyl)- 5,6-diphenyl-2-pyrazinecarboxamide;

25 *N*-(1-piperidinyl)- 5,6-bis(4-bromophenyl)-2-pyrazinecarboxamide;

- N*-(1-piperidiny)- 5,6-bis(4-methylphenyl)- 2-pyrazinecarboxamide;
N-(1-piperidiny)- 5,6-bis(4-methoxyphenyl)- 2-pyrazinecarboxamide;
N-(1-piperidiny)- 5,6-bis(4-chlorophenyl)- 2-pyrazinecarboxamide;
N-(1-piperidiny)- 5,6-bis(2-chlorophenyl)- 2-pyrazinecarboxamide;
5 *N*-cyclohexyl-5,6-diphenyl-2-pyrazinecarboxamide;
N-cyclohexyl-5,6-bis(4-bromophenyl)-2-pyrazinecarboxamide;
N-cyclohexyl-5,6-bis(4-methylphenyl)-2-pyrazinecarboxamide;
N-cyclohexyl-5,6-bis(4-methoxyphenyl)-2-pyrazinecarboxamide;
N-cyclohexyl-5,6-bis(4-chlorophenyl)- 2-pyrazinecarboxamide;
10 *N*-cyclohexyl-5,6-bis(2-chlorophenyl)-2-pyrazinecarboxamide;
N,5,6-triphenyl-2-pyrazinecarboxamide;
N-phenyl-5,6-bis(4-methylphenyl)-2-pyrazinecarboxamide;
N-phenyl-5,6-bis(4-methoxyphenyl)-2-pyrazinecarboxamide;
N-phenyl-5,6-bis(4-chlorophenyl)-2-pyrazinecarboxamide;
15 *N*-phenyl-5,6-bis(2-chlorophenyl)-2-pyrazinecarboxamide;
N-(1-piperidiny)- 5-(4-chlorophenyl)-6-(2,4-dichlorophenyl)-2-pyrazinecarboxamide; and
N-(1-piperidiny)- 6-(4-chlorophenyl)-5-(2,4-dichlorophenyl)-2-pyrazinecarboxamide;
and where applicable, optical isomers, tautomers, stereoisomers and racemates thereof as
well as pharmaceutically acceptable salts, solvates and crystalline forms thereof.
- 20
11. A compound of formula I as claimed in any previous claim for use as a medicament.
12. A pharmaceutical formulation comprising a compound of formula I, as defined in any
one of claims 1 to 10 and a pharmaceutically acceptable adjuvant, diluent or carrier.
- 25
13. Use of a compound of formula I



I

and pharmaceutically acceptable salts, prodrugs, solvates and crystalline forms thereof, in which

R^1 and R^2 independently represent :

- 5 a C_{1-6} alkyl group;
- an (amino) C_{1-4} alkyl- group in which the amino is optionally substituted by one or more C_{1-3} alkyl groups;
- an optionally substituted non-aromatic C_{3-15} carbocyclic group;
- a $(C_{3-12}$ cycloalkyl) C_{1-3} alkyl- group;
- 10 a group $-(CH_2)_r(phenyl)_s$, in which r is 0, 1, 2, 3 or 4, s is 1 when r is 0 otherwise s is 1 or 2 and the phenyl groups are optionally independently substituted by one, two or three groups represented by Z ;
- naphthyl;
- anthracenyl;
- 15 a saturated 5 to 8 membered heterocyclic group containing one nitrogen and optionally one of the following : oxygen, sulphur or an additional nitrogen wherein the heterocyclic group is optionally substituted by one or more C_{1-3} alkyl groups, hydroxy or benzyl ;
- 1-adamantylmethyl;
- a group $-(CH_2)_t$ Het in which t is 0, 1, 2, 3 or 4, and the alkylene chain is optionally
- 20 substituted by one or more C_{1-3} alkyl groups and Het represents an aromatic heterocycle optionally substituted by one, two or three groups selected from a C_{1-5} alkyl group, a C_{1-5} alkoxy group or halo;
- or R^1 represents H and R^2 is as defined above;
- or R^1 and R^2 together with the nitrogen atom to which they are attached represent a
- 25 saturated 5 to 8 membered heterocyclic group containing one nitrogen and optionally one

of the following : oxygen, sulphur or an additional nitrogen; wherein the heterocyclic group is optionally substituted by one or more C₁₋₃alkyl groups, hydroxy or benzyl ;

X is CO or SO₂ ;

5

Y is absent or represents NH optionally substituted by a C₁₋₃alkyl group;

R³ and R⁴ independently represent phenyl, thienyl or pyridyl each of which is optionally substituted by one , two or three groups represented by Z;

10

Z represents a C₁₋₃alkyl group, a C₁₋₃alkoxy group, hydroxy, halo, trifluoromethyl, trifluoromethylthio, trifluoromethoxy, trifluoromethylsulphonyl, nitro, amino, mono or di C₁₋₃alkylamino, mono or di C₁₋₃alkylamido, C₁₋₃alkylsulphonyl, C₁₋₃alkoxycarbonyl, carboxy, cyano, carbamoyl, mono or di C₁₋₃alkyl carbamoyl, sulphamoyl and acetyl; and

15

R⁵ is H, a C₁₋₃alkyl group, a C₁₋₃alkoxymethyl group, trifluoromethyl, a hydroxyC₁₋₃alkyl group, C₁₋₃alkoxycarbonyl, carboxy, cyano, carbamoyl, mono or di C₁₋₃alkylcarbamoyl, acetyl, or hydrazinocarbonyl of formula -CONHNR^aR^b wherein R^a and R^b are as previously defined for R¹ and R² respectively;

20

in the preparation of a medicament for the treatment or prophylaxis of obesity, psychiatric disorders such as psychotic disorders, schizophrenia and bipolar disorders, anxiety, anxio-depressive disorders, depression, cognitive disorders, memory disorders, obsessive-compulsive disorders, anorexia, bulimia, attention disorders, epilepsy, and related conditions, and neurological disorders such as dementia, neurological disorders,


25

Parkinson's Disease, Huntington's Chorea and Alzheimer's Disease, immune, cardiovascular, reproductive and endocrine disorders, septic shock, diseases related to the respiratory and gastrointestinal systems, and extended abuse, addiction and/or relapse indications.

30

14. A method of treating obesity, psychiatric disorders, psychotic disorders, schizophrenia and bipolar disorders, anxiety, anxio-depressive disorders, depression, cognitive disorders,

memory disorders, obsessive-compulsive disorders, anorexia, bulimia, attention disorders, epilepsy, and related conditions, neurological disorders, neurological disorders, Parkinson's Disease, Huntington's Chorea and Alzheimer's Disease, immune, cardiovascular, reproductive and endocrine disorders, septic shock, diseases related to the respiratory and gastrointestinal system, and extended abuse, addiction and/or relapse indications, comprising administering a pharmacologically effective amount of a compound of formula I



I

and pharmaceutically acceptable salts, prodrugs, solvates and crystalline forms thereof, in which

R^1 and R^2 independently represent :

a C_{1-6} alkyl group;

an (amino) C_{1-4} alkyl- group in which the amino is optionally substituted by one or more

C_{1-3} alkyl groups;

an optionally substituted non-aromatic C_{3-15} carbocyclic group;

a (C_{3-12} cycloalkyl) C_{1-3} alkyl- group;

a group $-(\text{CH}_2)_r(\text{phenyl})_s$, in which r is 0, 1, 2, 3 or 4, s is 1 when r is 0 otherwise s is 1 or 2 and the phenyl groups are optionally independently substituted by one, two or three groups

represented by Z;

naphthyl;

anthracenyl;

a saturated 5 to 8 membered heterocyclic group containing one nitrogen and optionally one of the following : oxygen, sulphur or an additional nitrogen wherein the heterocyclic

group is optionally substituted by one or more C_{1-3} alkyl groups, hydroxy or benzyl ;

1-adamantylmethyl;

a group $-(CH_2)_t$ Het in which t is 0, 1, 2, 3 or 4, and the alkylene chain is optionally substituted by one or more C_{1-3} alkyl groups and Het represents an aromatic heterocycle optionally substituted by one, two or three groups selected from a C_{1-5} alkyl group, a C_{1-5} alkoxy group or halo;

5 or R^1 represents H and R^2 is as defined above;

or R^1 and R^2 together with the nitrogen atom to which they are attached represent a saturated 5 to 8 membered heterocyclic group containing one nitrogen and optionally one of the following : oxygen, sulphur or an additional nitrogen; wherein the heterocyclic group is optionally substituted by one or more C_{1-3} alkyl groups, hydroxy or benzyl ;

10

X is CO or SO_2 ;

Y is absent or represents NH optionally substituted by a C_{1-3} alkyl group;

15 R^3 and R^4 independently represent phenyl, thienyl or pyridyl each of which is optionally substituted by one , two or three groups represented by Z;

Z represents a C_{1-3} alkyl group, a C_{1-3} alkoxy group, hydroxy, halo, trifluoromethyl, trifluoromethylthio, trifluoromethoxy, trifluoromethylsulphonyl, nitro, amino, mono or di C_{1-3} alkylamino, mono or di C_{1-3} alkylamido, C_{1-3} alkylsulphonyl, C_{1-3} alkoxycarbonyl, 20 carboxy, cyano, carbamoyl, mono or di C_{1-3} alkyl carbamoyl, sulphamoyl and acetyl; and

R^5 is H, a C_{1-3} alkyl group, a C_{1-3} alkoxymethyl group, trifluoromethyl, a hydroxy C_{1-3} alkyl group, C_{1-3} alkoxycarbonyl, carboxy, cyano, carbamoyl, mono or di C_{1-3} alkylcarbamoyl, 25 acetyl, or hydrazinocarbonyl of formula $-CONHN^aR^b$ wherein R^a and R^b are as previously defined for R^1 and R^2 respectively;


to a patient in need thereof.

15. A compound as defined in any one of claims 1 to 10 for use in the treatment of obesity.

30


16. A process for the preparation of a compound of formula I comprising

a) reacting a compound of formula II




II

5 in which R^3 , R^4 and R^5 are as previously defined with an amine of formula III




10 in an inert solvent in the presence of a coupling agent and optionally in the presence of a catalyst at a temperature in the range of -25°C to 150°C to give a compound of formula I in which X is CO; or

b) reacting a compound of formula IV




IV

15 in which R^3 , R^4 and R^5 are as previously defined and A represents halo with an amine of formula V



20 in an inert solvent and optionally in the presence of a catalyst at a temperature in the range of -25°C to 150°C to give a compound of formula I in which X is SO_2 .

17. A compound of formula II



II

- 5 in which R³, R⁴ and R⁵ are as previously defined with the exception of 5,6-diphenyl-2-pyrazinecarboxylic acid and 5,6-bis(4-methoxyphenyl)-2-pyrazinecarboxylic acid.

International Application No
PCT/GB 02/05742

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D241/24 A61K31/495 A61P25/18

B. FIELDS SEARCHED

IPC 7 C07D

EPO-Internal. WPI Data, CHEM ABS Data, BEILSTEIN Data

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	--	-----------------------

17

1-12

1-16

-/-

Y Patent family members are listed in annex.

*& document member of the same patent family

15/04/2003

Menegaki, F

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 02/05742

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>HOWLETT A C ET AL: "AZIDO- AND ISOTHIOCYANATO-SUBSTITUTED ARYL PYRAZOLES BIND COVALENTLY TO THE CB1 CANNABINOID RECEPTOR AND IMPAIR SIGNAL TRANSDUCTION" JOURNAL OF NEUROCHEMISTRY, NEW YORK, NY, US, vol. 74, no. 5, 2000, pages 2174-2181, XP001097394 ISSN: 0022-3042 figure 1</p>	1-16
A	<p>EP 0 397 859 A (TERUMO CORP) 22 November 1990 (1990-11-22) example 14; table 1</p>	17

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 02/05742

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
CH 458361	A	30-06-1968	NONE	
WO 9202513	A	20-02-1992	WO 9202513 A1	20-02-1992
			JP 6501926 T	03-03-1994
EP 0656354	A	07-06-1995	FR 2713224 A1	09-06-1995
			FR 2713225 A1	09-06-1995
			AT 154012 T	15-06-1997
			AU 685518 B2	22-01-1998
			AU 7899994 A	15-06-1995
			BR 1100984 A3	14-03-2000
			CA 2136893 A1	21-06-1995
			CN 1110968 A ,B	01-11-1995
			CZ 9403016 A3	14-06-1995
			DE 69403614 D1	10-07-1997
			DE 69403614 T2	22-01-1998
			DK 656354 T3	29-12-1997
			EP 0656354 A1	07-06-1995
			ES 2105575 T3	16-10-1997
			FI 945690 A	03-06-1995
			GR 3024470 T3	28-11-1997
			HK 1000599 A1	09-04-1998
			HU 71498 A2	28-11-1995
			IL 111719 A	28-10-1999
			JP 3137222 B2	19-02-2001
			JP 7309841 A	28-11-1995
			JP 2001026541 A	30-01-2001
			NO 944625 A	06-06-1995
			NZ 270025 A	26-09-1995
			PL 306067 A1	12-06-1995
			RU 2141479 C1	20-11-1999
			SG 68570 A1	20-06-2000
			SI 656354 T1	31-10-1997
			US 5624941 A	29-04-1997
			ZA 9409342 A	09-10-1995
EP 0397859	A	22-11-1990	JP 1128971 A	22-05-1989
			JP 1128972 A	22-05-1989
			JP 1824748 C	10-02-1994
			JP 5036435 B	31-05-1993
			JP 1135775 A	29-05-1989
			JP 1824749 C	10-02-1994
			JP 5036434 B	31-05-1993
			EP 0397859 A1	22-11-1990
			WO 8904308 A1	18-05-1989

THIS PAGE BLANK (USPTO)